

NIPP: Nursing Home Infection Prevention Program...*Nipping Infections in the Bud*[©]
Surveillance Criteria for Urinary Tract Infection With Indwelling Urinary Catheter*
Per the CDC National Healthcare Safety Network Surveillance Protocol for LTCF

*Indwelling urinary catheter in place or removed within 2 calendar days prior to event onset (catheter removed = day 1)

Does the patient have one or more of the following signs and symptoms?

- Fever (Single temp $\geq 37.8^{\circ}\text{C}$ or $>100^{\circ}\text{F}$ or $>37.2^{\circ}\text{C}$ or 99°F on repeated occasions or an increase of $>1.1^{\circ}\text{C}$ or $>2^{\circ}\text{F}$ over baseline)
- Rigors
- New onset hypotension, with no alternate non-infectious cause
- New onset confusion/functional decline with no alternate diagnosis AND leukocytosis (>10000 cells/ mm^3 or Left shift $> 6\%$ or 1500 bands/ mm^3)
- New or marked increase in suprapubic tenderness
- New or marked increase in costovertebral angle pain or tenderness
- Acute pain, swelling, or tenderness of the testes, epididymis, or prostate
- Purulent discharge from around the catheter insertion site



- Does the patient have a positive urine culture?**
- Specimen collected from indwelling catheter or clean catch (catheter removed) voided urine yields no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml
- OR
- Specimen collected from in/out straight catheter (indwelling catheter removed) and yields no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml

